



**PCRBIO SYSTEMS**  
simplifying research

## PCRBIO HS VeriFi™ Mix Red

[www.pcrbio.com](http://www.pcrbio.com)

**\*\* Special pack for Weizmann Institute of Science \*\***

**2x PCRBIO HS VeriFi™ Mix Red, 1 x 1.25ml (50x50µl rxns / 250x20µl rxns)**

### Product description:

PCRBIO HS VeriFi™ Mix Red is a convenient high fidelity mix with AptaLock™ hot start technology for highly precise PCR. This 2x ready mix is designed for PCR applications where greater sequence accuracy is required, together with improved PCR success rates of long and challenging templates. The mix contains a red dye suitable for direct loading and tracking during agarose gel electrophoresis.

PCRBIO HS VeriFi™ Mix Red contains the highly processive PCRBIO HS VeriFi™ Polymerase, developed for robust and versatile high fidelity PCR. The enzyme is derived from Pfu DNA polymerase for its 3'-5' exonuclease (proofreading) activity. Several proprietary mutations significantly improve DNA binding and processivity, resulting in shorter extension times (30s/kb), higher yields and the ability to amplify longer and more difficult targets, including eukaryotic genomic templates in excess of 17.5kb.

PCRBIO's innovative AptaLock™ technology uses a proprietary aptamer-like molecule that reversibly inhibits both the 3'-5' exonuclease activity and 5'-3' polymerase activity of the enzyme at ambient temperatures. This unique hot start molecule prevents primer dimer formation and non-specific amplification to maximise the sensitivity and specificity of your PCR. This feature makes the enzyme highly suitable for multiplexing and enables reactions to be set up at room temperature.

The enhanced accuracy of PCRBIO HS VeriFi™ Polymerase results in fidelity that is approximately 100 times higher than Taq DNA polymerase, making it ideal for applications such as cloning, site-directed mutagenesis and sequencing.

Component	100 x 50µL rxns	500 x 50µL rxns
2x PCRBIO HS VeriFi™ Mix Red	2 x 1.25mL	10 x 1.25mL

PCRBIO HS VeriFi™ Mix Red uses an advanced buffer system including dNTPs, Mg and enhancers, enabling high fidelity PCR of a wide range of targets and fragment sizes regardless of GC content. PCR products generated with this range of products are blunt ended.

### Shipping and storage

On arrival the kit should be stored between -30°C and -15°C. If stored correctly the kit will retain full activity for 12 months. The kit can be stored at 4°C for 1 month.

### Limitations of product use

The product may be used for in vitro research purposes only.

### Technical support

Help and support is available on our website at <https://pcrbio.com/resources/> including answers to frequently asked technical questions. For technical support and troubleshooting you can submit a technical enquiry online, or alternatively email [technical@pcrbio.com](mailto:technical@pcrbio.com) with the following information:

- Amplicon size
- Reaction setup
- Cycling conditions
- Screen grabs of gel images

## Important considerations

**2x PCR BIO HS VeriFi™ Mix Red:** The 2x mix contains PCR BIO HS VeriFi™ Polymerase, 6mM MgCl<sub>2</sub>, 2mM dNTPs, enhancers, stabilizers, and a red dye for tracking during agarose electrophoresis. It is not recommended to add further PCR enhancers or MgCl<sub>2</sub> to the reaction. The mix composition has been optimised to maximise PCR success rates.

**Primers:** Primers should have a predicted melting temperature of around 60°C, using default Primer 3 settings (<http://bioinfo.ut.ee/primer3/>). The final primer concentration in the reaction should be between 0.2µM and 0.6µM.

**Denaturation:** Denaturation should be performed at 95°C. However, if the presence of high GC regions results in low yields, increasing the melting temperature to 98-100°C can improve the amount of product.

**Annealing:** We recommend performing a temperature gradient to experimentally determine the optimal annealing temperature. Alternatively, we recommend a 60°C annealing temperature then increase in 2°C increments if non-specific products are present.

**Extension:** Optimal extension is achieved at 72°C. The optimal extension time is dependent on amplicon length and complexity of template. 30 seconds per kilobase (kb) is recommended for most applications. Two-step cycling protocols may also be used with combined annealing and extension at 68-75°C.

**Multiplex PCR:** The optimal extension time for multiplex reactions will be dependent on the complexity of template, the length of amplicons, and the number of targets. We recommend starting with the extension time of the longest fragment, and then increasing in increments of between 10 and 30 seconds if necessary.

**Agarose gel electrophoresis dye migration:** The 2x mix contains a red dye for tracking during agarose gel electrophoresis. In a 2% agarose TAE gel the dye migrates at a rate equivalent to 350bp of DNA. In a 1% agarose TAE gel the dye migration rate is equivalent to 600bp of DNA.

## Reaction setup

1. Prepare a master mix on ice based on the following table:

Reagent	25µL reaction	50µL reaction	Final concentration	Notes
2x PCR BIO HS VeriFi™ Mix Red	12.5µL	25.0µL	1x	
Forward primer (10µM)	1.0µL	2.0µL	400nM	See above for optimal primer design
Reverse primer (10µM)	1.0µL	2.0µL	400nM	
Template DNA	<100ng genomic DNA <5ng less complex DNA	<200ng genomic DNA <10ng less complex DNA	variable	
PCR grade dH <sub>2</sub> O	Up to 25µL final volume	Up to 50µL final volume		

2. Cycle using conditions based on the following table:

Cycles	Temperature	Time	Notes
1	95°C	1min	Initial denaturation
25-35	95°C	15 seconds	Denaturation (see above for high GC templates)
	60°C to 75°C	15 seconds	Anneal
	72°C	30 seconds / kb	Extension (see above for multiplex PCR)